

## PESTICIDE COMPOSITIONS

This invention is related to the field of compositions useful in the control of pests that eat cellulose, such as, termites.

Cellulose is the most abundant composition on the planet. It is a  
5 polymer that is made from glucose. Wood contains 50 weight percent cellulose and cotton contains 90 weight percent cellulose. Cellulose is a generic term for a composition that contains alpha cellulose, beta cellulose, and gamma cellulose. Alpha cellulose has a much higher degree of polymerization ("DP") than beta or gamma cellulose. Alpha Cellulose has a  
10 DP in the thousands depending on the source of the alpha cellulose. Alpha cellulose can be made into microcrystalline cellulose. Microcrystalline cellulose has a DP of less than 400.

Pests that eat cellulose are very destructive. Termites are very destructive easily causing billions of dollars in damage each year.  
15 Research is constantly being conducted to find new measures that can better control termites, especially termite species that have been difficult to control in the past.

This invention provides a solution to the problems of controlling pests that eat cellulose.

20 This invention is related to the field of compositions useful in the control of pests that eat cellulose, such as, termites, where said

composition is compacted and where said composition comprises alpha-cellulose, water, and a pesticide.

US 2003/0152605 A1 discloses an optimum density termite composition. An optimum density was required because the applicants  
5 wanted more cellulose compacted into less volume thereby, according to the application, providing a longer term feeding composition. The composition comprises "purified cellulose" (it is not readily apparent from the application what this term means) or microcrystalline cellulose in a compacted form where the composition has a density greater than 1.033  
10 grams per cubic centimeter. It is apparent that the applicants are not disclosing any composition that contains water in any amount. This is especially apparent when considering their problems with powder cellulose as disclosed in paragraph 0014 and their solution for deaerating the powder in paragraph 019.

## 15 **PESTS**

Any cellulose eating pest can be controlled by the inventive compositions disclosed herein. In particular, termites can be controlled by the inventive compositions herein, especially subterranean termites such as *Reticulitermes* spp. and *Coptotermes* spp. Suitable examples of  
20 termites that can be controlled are: *Reticulitermes flavipes*; *Reticulitermes virginicus*; *Reticulitermes Hesperus*; *Heterotermes aureus*; *Coptotermes formosanus*; *Reticulitermes speratus*; *Reticulitermes grassei*;

Reticulitermes santonensis; Macrotermes gilvus; and Reticulitermes hageni.

### ALPHA CELLULOSE

Alpha cellulose is readily available. It can be purchased from a variety of sources. One particular source is International Fiber Corporation. It is particularly preferred if the alpha cellulose is in powdered form.

### PESTICIDES

Examples of suitable insecticides that may be used are:

(a) Pyrethroids, such as permethrin, cypemethrin, fenvalerate, esfenvalerate, deltamethrin, cyhalothrin, lambda-cyhalothrin, gamma-cyhalothrin, bifenthrin, fenpropathrin, cyfluthrin, tefluthrin, fish safe pyrethroids (for example ethofenprox), natural pyrethrin, tetramethrin, s-bioallethrin, fenfluthrin, prallethrin, 5-benzyl-3-furylmethyl-(E)-(1R,3S)-2,2-dimethyl-3-(2-oxothiolan-3-ylidenemethyl) cyclopropane carboxylate, or any of their insecticidally active isomers;

(b) Organophosphates, such as, methidathion, chlorpyrifos-methyl, profenofos, sulprofos, acephate, methyl parathion, azinphos-methyl, demeton-s-methyl, heptenophos, thiometon, fenamiphos, monocrotophos, profenofos, triazophos, methamidophos, dimethoate, phosphamidon, malathion, chlorpyrifos, chlorpyrifos-methyl, phosalone, terbufos, fensulfothion, fonofos, phorate, phoxim, pirimiphos-methyl, pirimiphos-ethyl, fenitrothion, fosthiazate or diazinon;

- (c) Carbamates (including aryl carbamates), such as fenoxycarb, alanycarb, pirimicarb, triazamate, cloethocarb, carbofuran, furathiocarb, ethiofencarb, aldicarb, thiofurox, carbosulfan, bendiocarb, fenobucarb, propoxur, methomyl or oxamyl;
- 5 (d) Benzoyl ureas, such as lufenuron, novaluron, noviflumuron, teflubenzuron, diflubenzuron, triflumuron, hexaflumuron, flufenoxuron or chlorfluazuron;
- (e) Organic tin compounds, such as cyhexatin, fenbutatin oxide or azocyclotin;
- 10 (f) Pyrazoles, such as tolfenpyrad, pyridaben, tebufenpyrad and fenpyroximate;
- (g) Macrolides, such as avermectins or milbemycins, for example abamectin, emamectin benzoate, ivermectin, milbemycin, spinosad or azadirachtin;
- 15 (h) Hormones or pheromones;
- (i) Organochlorine compounds such as endosulfan, benzene hexachloride, DDT, chlordane or dieldrin;
- (j) Amidines, such as chlordimeform or amitraz;
- (k) Chloronicotinyl compounds such as diofenolan, clothianidin,
- 20 thiacloprid, imidacloprid, thiacloprid, acetamiprid, nitenpyram or thiamethoxam;

(l) Diacylhydrazines, such as halofenozide, tebufenozide, chromafenozide or methoxyfenozide;

(m) Diphenyl ethers, such as diofenolan or pyriproxifen;

(n) Indoxacarb;

5 (o) Chlorfenapyr;

(p) Pymetrozine;

(q) Diafenthiuron;

(r) Toxins of microbial origin such as *B.acillus thuringiensis* endo- or exotoxins;

10 (s) Phenylpyrazoles such as fipronil, vanilliprole, etiprole or acetoprole;

(t) Pyridalyl; or

(v) hydramethylnon

In addition to the major pesticides listed above, other pesticides  
15 having particular targets may be employed if appropriate for the intended utility of the inventive composition. Alternatively, insecticides, or acaricides, specific for particular insect species/stages may also be included in the inventive compositions (for example acaricidal ovo-larvicides, such as clofentezine, flubenzimine, hexythiazox or tetradifon; acaricidal motilicides,  
20 such as dicofol or propargite; acaricides, such as acequinocyl, fenazaquin, spiroticlofen, etoxazole, bromopropylate or chlorobenzilate; or growth

regulators, such as cyromazine, methoprene, chlorfluazuron or diflubenzuron).

Examples of suitable insecticide synergists that may be used as a further active ingredient in the inventive compositions include piperonyl  
5 butoxide, sesamex, safroxan and dodecyl imidazole.

Specific examples of preferred pesticides are thiamethoxam, abamectin, emamectin benzoate, spinosad, chlorpyrifos, chlorpyrifos-methyl, profenofos, lufenuron, indoxacarb, lambda-cyhalothrin, pymetrozine, pirimicarb, methidathion, imidacloprid, acetamiprid,  
10 thiacloprid, fipronil, methoxyfenozide, chlorfenapyr, pyridaben, novaluron, noviflumuron, pyridalyl, propargite and piperonyl butoxide.

Mixtures of pesticides are also useful and many of the above can be synergistically used together.

However, it is most preferred to use a slow acting pesticide,  
15 especially with termites, so that the termites can take the pesticide, or pesticides, back to their colony and poison other colony members.

#### **PREPARATION AND USE**

In general the amount of alpha cellulose and water to use is not critical and can vary by a wide amount depending on the other  
20 components (such as pesticide(s), binder(s), attractant(s), etc.) added to mixture of alpha cellulose and water. It is preferred that the mixture of alpha cellulose, water, and the other components, if any, upon compacting,

have a density greater than 1 gram per cubic centimeter. Densities less than 1 gram per cubic centimeter can be used but are not preferable in most cases. Suitable ranges for alpha cellulose and water are in Table 1.

Component	Table 1: Weight Percent (based on total weight of both components)		
	Preferred Range	More Preferred Range	Most Preferred Range
Alpha Cellulose	60 to 99	65 to 95	70 to 90
Water	40 to 1	35 to 5	30 to 10

5 In general the amount of pesticide to use is also not critical.

Amounts from 0.0001 to 10 weight percent based on the weight of the inventive composition can be used (that is, alpha cellulose, water, pesticide, plus any other components desired).

The compacted composition of alpha cellulose, water, pesticide,  
10 plus any other components desired can take any useful form. A useful form is a form that the desired pest can eat. Such forms are tablets, briquettes, pellets, granules, etc. These types of forms can be made by any process known in the art. In general, more water is needed to form good pellets and less water is needed to form good briquettes.

15 In an unexpected and surprising manner, it has been discovered that termites that have in the past been difficult to control with baits, feed especially well upon briquettes as opposed to pellets or tablets, possibly because the briquettes of this invention can absorb liquids more easily. Consequently, it is preferred to compact the inventive composition in the

shape of a briquette. One machine that can do this process is Komarek model B-100-A two roll mill available from K. R. Komarek Inc., 1825 Estes Ave., Elk Grove Village, IL 60007.

Once the inventive composition has been compacted, such as into  
5 briquettes, it is preferred to dry the briquettes. This drying can occur in any manner known in the art that will remove a portion of the water. This may seem paradoxical, in that it is desirable to have water in the compacted composition. After all, water is desired by cellulose eating pests, especially termites. However, while water is removed from the compacted  
10 composition it is not entirely removed from the compacted composition. At least a portion of the water that is removed from the compacted composition leaves voids in the compacted composition.

The compacted composition can then be used as a bait to control cellulose eating pests, such as subterranean termites. Many methods are  
15 known to bait such pests. In a preferred embodiment of the invention the compacted composition is placed in the ground, perhaps inside another tube that allows access for termites, and prior to placing the compacted composition in the ground or tube, or leaving it in the ground or tube, additional liquid (such as water, or a mixture of water and sugar, or a  
20 mixture of water, sugar, and salts) is contacted with the compacted composition. This method can attract even more termites to the now moistened, compacted composition, causing more feeding and more



pesticide delivery to the termites and the colony. Since the compacted composition was first dried before being moisten the liquid can readily fill any voids left behind by the drying of the compacted composition.

Optional ingredients to include in the inventive composition include,  
5 but are not limited to, a preservative to retard fungal growth, and a protectant such as a bittering agent to provide a safety factor for exposed bait.

An attractant is defined as any substance or combination of substances which will lure pests. Examples of attractants are carbon  
10 dioxide and terpenes.

Feeding stimulants that can be used in the inventive composition are, for example, sugar, such as powdered sucrose, high fructose corn syrup, polyhydroxy alcohols such as glycerin, and starch.

Examples of preservatives useful in the present invention are 1,2-  
15 benzisothiazolin-3-one (PROXEL GXL<sup>®</sup>., Avecia Inc. Wilmington, Del. 19850) methyl paraben (p-hydroxybenzoic acid methyl ester) and propyl paraben (n-propyl p-hydroxybenzoate). Other known fungistats would also be effective in increasing the longevity of the bait and retarding mold growth.

20 The headings used herein are meant to be as a guide and are not to be used to interpret the scope of the invention.

## EXAMPLES

These examples are provided to illustrate the invention. They are not to be used for limiting the scope of the invention.

### 5 Example A: Preparation of Pesticide Concentrate

Pesticide Concentrate Table	
Wt %	Ingredient
50.5	Noviflumuron
38.1	Water
10.4	Pluronic P-104
0.7	Proxel GXL
0.3	Antifoam B

A pesticide concentrate containing the amounts of materials in the pesticide concentrate table was prepared as follows.

Pluronic P-104 and water were mixed together to form a solution  
10 containing 23.3 weight percent Pluronic P-104 based on the total weight of the mixture ("First Mixture"). The Pesticide concentrate was made by mixing together the First Mixture, the noviflumuron, the Proxel GXL, and the Antifoam B in the amounts required to achieve the indicated weight percents.

15

### Example 1-A: Preparation of a Preferred Embodiment

In this example, a preferred embodiment of the inventive compositions disclosed herein is made.

Ninety kg of cellulose (BH-100 from International Fiber Corporation) was weighed into a blender. A first mixture of pesticide concentrate (see Example A) with water (1 part pesticide concentrate: 3 parts water) was then sprayed into the blender. Upon mixing, this produced a second  
5 mixture containing 0.5 weight percent pesticide, 9.2 weight percent water, and 90.3 weight percent cellulose, based on the weight of the second mixture.

Additional water was sprayed into the blender to produce a third mixture that could be compacted into briquettes. The third mixture  
10 contained 0.4 weight percent pesticide, 27.9 weight percent water, and 71.7 weight percent cellulose, based on the weight of the third mixture.

The third mixture was then fed into a compaction device. The compaction device was the Komarek briquetter, model B-100-A two-roll machine. The machine was run under the following conditions to produce  
15 briquettes: a compaction force of 1,300 kg/cm<sup>2</sup>; a feed rate of 400-500 g/min; and a tip speed of 5 feet per minute.

The briquettes coming off the compaction device were then screened with a vibratory screener. The material that passed through a 4-mesh screen (4,750 micron size) was collected for recycling to the feed.

20 The briquettes were then dried to remove a portion of the water.

**Example 1-B: Preparation of another Preferred Embodiment**

In this example, a preferred embodiment of the inventive compositions disclosed herein is made.

68.18 kg of cellulose (BH-100 from International Fiber Corporation)  
5 was weighed into a blender. A first mixture of pesticide concentrate (see  
Example A) with water (0.698 kg pesticide concentrate and 2.116 kg  
water) was then mixed into the blender. An additional 24.3 kg of water was  
then added to the blender. After thorough mixing, the mixture was fed into  
a model CL-2 California Pellet mill to produce 3/16" diameter pellets. 500  
10 – 600 grams per minute of wet pellets were produced at a rotational speed  
of 620 feet per min. The material coming off the pellet mill was screened.  
Product that passed through a 4-mesh screen (4,750  $\mu$ ) was disposed.  
The pellets were dried to remove a portion of the water.

**15 Example 1-C: Preparation of another Preferred Embodiment**

In this example, a preferred embodiment of the inventive compositions disclosed herein is made.

90.91 kg of cellulose (BH-100 from International Fiber Corporation)  
was weighed into a blender. 16.243 kg of BF-20 starch (From Grain  
20 Processing Corporation) was then mixed into the blender. A first mixture of  
pesticide concentrate (see Example A) with water (1.1 kg pesticide  
concentrate and 3.3 kg water) was then mixed into the blender. An

additional 30.419 kg of water was then added to the blender. After thorough mixing, the mixture was fed into a model CL-2 California Pellet mill to produce 3/16" diameter pellets. 400 – 600 grams per minute of wet pellets were produced at a rotational speed of 620 feet per min. The material coming off the pellet mill was screened. Product that passed through a 4-mesh screen (4,750  $\mu$ ) was disposed. The pellets were dried to remove a portion of the water.

#### **Example 2: Importance of Moist Baits**

10           This example shows the importance of moisture to termites.

          The moist bait was prepared in accordance with Example 1-B. Additional water was added to the bait as indicated (either 0.43 ml/g; 0.85 ml/g; Or 2 ml/g; water/gram of bait)

          The dry bait was prepared in accordance with Example 1-B.

15           However no additional water was added.

          A one-way pair choice testing environment was used. The termites were placed in a harborage and a tube allowed them a passage to the food environment. In the food environment, the termites were given a choice of two different food sources. The termites were allowed to feed on the food choices for 7 days. Between 4 and 12 reps were performed and each rep had 100-200 termite workers. Bait consumption was measured as the difference between the pre-weight and the post weight.

At 0.43 ml/g *R. flavipes*, *R. speratus*, *C. gestrol/travians*, and *H. aureus* preferred the moist bait over the dry bait. The only other species tested had no preference *R. hesperus*.

At 0.85 ml/g *R. speratus*, *C. gestrol/travians*, *H. aureus*, *R. grassei*,  
5 *C. formosanus*, and *R. virginicus* preferred the moist bait over the dry bait.  
*R. flavipes* and *R. hesperus* showed no preference.

At 2 ml/g *R. speratus*, and *C. gestrol/travians* preferred the moist bait over the dry bait. *R. flavipes*, *R. hesperus*, *H. aureus* showed no preference. *R. grassei* actually preferred the dry bait over the moist bait.

10 This example shows the importance of moisture to termite feeding, and hence to the importance of the inventive composition's ability to absorb moisture.